

**Infants and Toddlers Who Require Specialty Services and Supports**  
**(birth to 47 months of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+**

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis	
100	<b>Post traumatic Stress Disorder</b>	308	Acute reaction to stress** (requires 4 <sup>th</sup> digit subclassification listed below). Includes: • Catastrophic stress • Gross stress reaction (acute) • Transient disorders in response to exceptional physical or mental stress which usually subside within hours or days	308.3	Acute Stress Disorder
		308.0	Predominant disturbance of emotions • Anxiety as acute reaction to exceptional (gross) stress		
		308.1	Predominant disturbance of consciousness Fugues as acute reaction to exceptional (gross) stress		
		308.2	Predominant psychomotor disturbance • Agitation states as acute reaction to exceptional (gross) stress • Stupor as acute reaction to exceptional (gross) stress		
		308.3	Other acute reactions to stress** • Acute situational disturbance • Brief or acute posttraumatic stress disorder		
		308.4	Mixed disorders as reaction to stress		
		308.9	Unspecified acute reaction to stress		
		309.81	Prolonged posttraumatic stress disorder** • Chronic posttraumatic stress disorder	309.81	Post traumatic Stress Disorder
150	<b>Deprivation/Maltreatment Disorder</b>	313.89	Emotional Disturbance of Childhood NEC**	313.89	Reactive Attachment Disorder of Infancy or Early Childhood
		313.9	Unspecified emotional Disturbance of Childhood or adolescence		

+ To be used as a guide in Clinical Assessment/Diagnosis

**Infants and Toddlers Who Require Specialty Services and Supports**  
**(birth to 47 months of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+**

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis	
200	<b>Disorder of Affect</b>				
210	Prolonged Bereavement/Grief Reaction	309.00	Adjustment disorder with depressed mood • Grief reaction	309.0	Adjustment Disorder with Depressed Mood
		309.1	Prolonged depressive reaction**		
220	Anxiety Disorder of Infancy & Early Children	300.00	Anxiety state, unspecified** • Anxiety: neurosis, reaction, state (neurotic) • Atypical anxiety disorder	300.00	Anxiety Disorder NOS
		300.01	Panic disorder. Panic: attack, state	300.01	Panic disorder without
		300.02	Generalized anxiety disorder	300.02	Generalized Anxiety Disorder
		309.21	Separation anxiety disorder	309.21	Separation Anxiety Disorder
		313.0	Overanxious disorder** • Anxiety and fearfulness of childhood and adolescence • Overanxious disorder of childhood and adolescence		
221	Separation Anxiety	309.21	Separation anxiety disorder	309.21	Separation Anxiety Disorder
222	Specific Phobia	300.29	Other isolated or specific phobias	300.29	Specific Phobia
223	Social Anxiety Disorder	300.23	Social Phobia	300.02	Social Phobia
		313.2	Sensitivity, shyness and social withdrawn disorder (requires 5 <sup>th</sup> digit subclassification—listed below)		
		313.21	Shyness disorder of childhood Sensitivity reaction of childhood or adolescence		
		313.22	Introverted disorder of childhood --Social withdrawal of childhood or adolescence --Withdrawal reaction of childhood or adolescence		
224	Generalized Anxiety Disorder			300.02	Generalized Anxiety Disorder
225	Anxiety Disorder NOS			300.0	Anxiety Disorder NOS

**Infants and Toddlers Who Require Specialty Services and Supports**  
**(birth to 47 months of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+**

<b>DC 0-3 R Diagnosis</b>		<b>ICD 9 CM Diagnosis</b> <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		<b>DSM IVR Diagnosis</b>	
230	<b>Depression of Infancy and Early Childhood</b>				
231	Type I Major Depression	296.2X	Major Depressive Disorder, Single Episode, Unspecified 5 <sup>th</sup> digit: 0 unspecified 1 mild, 2 moderate, 3 severe without mention of psychotic behavior, 4 severe with specified as with psychotic behavior, 5 in partial or unspecified remission, 6 in full remission	296.20	Major Depressive Disorder, Single Episode, Unspecified
		296.3X	Major Depressive Disorder, Recurrent, Unspecified 5 <sup>th</sup> digit: same as above	296.30	Major Depressive Disorder, Recurrent, Unspecified
232	Type II Depressive Disorder NOS	311	Depressive Disorder, not elsewhere classified --Depressive Disorder NOS --Depressive State NOS--Depression NOS	311	Depressive Disorder NOS
		313.1	Misery and unhappiness disorder		
		300.4	Neurotic Depression --Anxiety depression --Depression with anxiety --Depressive reaction --Dysthymic Disorder --Neurotic depressive State --Reactive Depression	300.4	Dysthymic Disorder
240	<b>Mixed Disorder of Emotional Expressiveness</b>	313.9	Unspecified emotional disturbances of childhood or adolescence	313.9	Disorder of Infancy, Childhood, or Adolescence NOS

**Infants and Toddlers Who Require Specialty Services and Supports**  
**(birth to 47 months of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+**

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis	
300	<b>Adjustment Disorder</b>	309	Adjustment Reaction (309.0—309.9)	309.9	Adjustment Disorder: Unspecified
		309.0	Brief depressive reaction --Adjustment disorder with depressed mood --Grief reaction		
		309.24	Adjustment reaction with anxious mood	309.24	Adjustment Disorder with Anxiety
		309.28	Adjustment reaction with mixed emotional features	309.28	Adjustment Disorder with Mixed Anxiety and Depressed Mood
		309.3	With predominant disturbance of conduct --conduct disturbance as adjustment reaction --destructiveness as adjustment reaction	309.3	Adjustment Disorder with Disturbance of Conduct
		309.4	With mixed disturbances of emotions and conduct	309.4	Adjustment Disorder with Mixed Disturbance of Emotions and Conduct
		309.29	Other --culture shock	309.9	Adjustment Disorder NOS
		309.82	Adjustment reaction with physical symptoms	309.9	Adjustment Disorder NOS
		309.83	Adjustment reaction with withdrawal --Elective mutism as adjustment reaction --Hospitalism (in children) NOS	313.23 309.9	Selective Mutism Adjustment Disorder NOS
		309.9	Adjustment Disorder NOS		
		309.9	Adjustment Disorder NOS		
		309.9	Adjustment Disorder NOS		
<b>400*– Regulation Disorders of Sensory Processing</b> Sensory Processing when coupled with parent/child interaction					
410*	Hypersensitive	312.9	Unspecified disturbance of conduct	312.9	Disruptive Behavior Disorder NOS
411*	Type A: Fearful/Cautious	313.0	Overanxious disorder** --Anxiety and fearfulness of childhood and adolescence --Overanxious disorder of childhood and	300.02	Generalized Anxiety Disorder

**Infants and Toddlers Who Require Specialty Services and Supports**  
**(birth to 47 months of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+**

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis	
			adolescence (Note: Also see 313.2, 313.21 or 313.22 for Sensitivity, shyness and social withdrawn disorder)		
412*	Type B: Negative/Defiant	312.81	Conduct Disorder—Child onset type (Note: for young child over 36 months of age)	312.81	Conduct Disorder—Child onset (Note: for young child over 36 months of age)
		312.9	Unspecified disturbance of conduct	312.9	Disruptive Behavior Disorder NOS
		313.81	Oppositional Defiant Disorder	313.81	Oppositional Defiant Disorder
420*	Hypo/Underresponsive	314.00	Attention deficit disorder Without mention of hyperactivity --Predominantly inattentive type	314.00	Attention Deficit Disorder— Predominately inattentive type
430*	Sensory Stimulation – Seeking/Impulsive	314.0	Attention Deficit Disorder --Requires 5 <sup>th</sup> digit subclassification-listed below	312.30 314.01	Impulse Control Disorder NOS ADHD-Hyperactive impulsive disorder
		314.01	With hyperactivity—combined type --Overactivity NOS --Predominantly hyperactive/impulsive		
		314.9	Unspecified hyperkinetic syndrome --Hyperkinetic reaction of childhood or adolescence NOS --Hyperkinetic syndrome NOS	314.9	
500*	<b>Sleep Disorder</b>				
510*	Sleep Onset Disorder	307.40	Nonorganic sleep disorder, unspecified	307.47	Dyssomnia NOS or Parasomnia NOS
		307.41	Transient disorder of initiating or maintaining sleep <ul style="list-style-type: none"> <li>• Hyposomnia associated with acute or intermittent emotional reactions or conflicts</li> <li>• Insomnia associated with acute or intermittent emotional reactions or conflicts</li> <li>• Sleeplessness associated with</li> </ul>	327.02	Insomnia Related to....

**Infants and Toddlers Who Require Specialty Services and Supports**  
**(birth to 47 months of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+**

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis	
		307.42	Persistent disorder of initiating or maintaining wakefulness • Hyposomnia, insomnia, or sleeplessness associated with: anxiety, conditioned arousal, depression (major) (minor), psychosis	307.42 or 327.02	Primary Insomnia Or Insomnia Related to...
		307.43	Transient disorder of initiating or maintaining wakefulness • Hypersomnia associated with acute or intermittent emotional reactions or conflicts	307.15	Hypersomnia Related to....
		307.44	Persistent disorder of initiating or maintaining wakefulness • Hypersomnia associated with depression (major) (minor)	307.44 or 327.15	Primary Hpersomnia or Hypersomnia Related to....
		307.45	Circadian Rhythm Sleep Disorder of nonorganic origin	307.30 or 327.31	Circadian Rhythm Sleep Disorder, Unspecified type or Delayed Sleep Phase Type
520*	Night –Waking Disorder	307.46	Somnambulism or night terrors	307.46	Sleep Terror Disorder or Sleepwalking Disorder
		307.47	Other dysfunctions of sleep stages or arousal from sleep Nightmares: NOS, REM-sleep type, sleep drunkenness	307.47	Dyssomnia NOS, Nightmare Disorder, or Parasomnia NOS
		307.48	Repetitive intrusions of sleep Repetitive intrusions of sleep with: atypical polysomnographic features, environmental disturbances, repeated REM-sleep interruptions	307.47	Parasomnia NOS
		307.49	Other “Short sleeper,” subjective insomnia complaint	307.42	Primary Insomnia
600	<b>Feeding Behavior Disorders</b>				
601	Feeding Behavior Disorders	307.50 307.59	Eating Disorder, unspecified Other • Infantile feeding disturbances of	307.59	Feeding Disorder of Infancy or Early Childhood

+ To be used as a guide in Clinical Assessment/Diagnosis

**Infants and Toddlers Who Require Specialty Services and Supports**  
**(birth to 47 months of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+**

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis	
			nonorganic origin • Loss of appetite—of nonorganic origin		
602	Feeding Disorders of Caregivers Caregiver/Infant Reciprocity	307.59	Other • Infantile feeding disturbances of nonorganic origin • Loss of appetite—of nonorganic origin	307.59	Feeding Disorder of Infancy or Early Childhood
603	Infantile Anorexia	307.59	Other • Infantile feeding disturbances of nonorganic origin • Loss of appetite—of nonorganic origin	307.59	Feeding Disorder of Infancy or Early Childhood
800	<b>Other Disorders</b>				

\* Not Axis I Diagnoses

References

American Psychiatric Association, Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR) Fourth Edition (Text Revision), 2000.

National Center for Health Statistics, International Classification of Diseases, Ninth Revision, Clinical Modification, 2004.

Zero to Three , Diagnostic Classification of Mental Health and Developmental Disorders of Infancy and Early Childhood: Revised Edition (DC: 0-3 R), Washington, DC, 2005.

Maine DC 0-3 Committee, Crosswalk for DC 0-3 to ICD-9 Diagnosis Coding. 2004.

Perez, L. and Newman, M., “Crosswalk” for San Mateo, California, Developmental Approach to Infant Mental Health Diagnosis.

In addition, the Crosswalk developed by Florida was used as a reference.

+ To be used as a guide in Clinical Assessment/Diagnosis